

	<b>Converting conventional radiographic examination data of trabecular bone pattern values into density measurement values using intraoral digital images</b>
Journal	Oral Radiology
Publisher	Springer Japan
ISSN	0911-6028 (Print) 1613-9674 (Online)
Issue	Volume 25, Number 2 / December, 2009
Category	Original Article
DOI	10.1007/s11282-009-0024-y
Pages	129-134
Subject Collection	Medicine
SpringerLink Date	Tuesday, September 15, 2009

**Menik Priaminiarti<sup>3</sup>**, **Budi Utomo<sup>1</sup>**, **R. Susworo<sup>2</sup>** and **Hanna Bachtiar Iskandar<sup>3</sup>**

- (1) Department of Population and Biostatistics, Faculty of Public Health, University of Indonesia, Jakarta, Indonesia
- (2) Department of Radiotherapy, Faculty of Medicine, University of Indonesia, Jakarta, Indonesia
- (3) Department of Dental and Maxillofacial Radiology, Faculty of Dentistry, University of Indonesia, Jl. Salemba Raya No.4, Jakarta, 10430, Indonesia

**Received:** 25 November 2008 **Accepted:** 31 July 2009 **Published online:** 9 September 2009

### **Abstract**

*Objectives* To determine the conversion value of grayscale density measurements from intraoral conventional radiographic examinations of the edentulous maxilla and mandible using intraoral digital radiography.

*Methods* Periapical radiography examinations with a pararelling technique, both conventional and digital, were performed on 18 male and 34 female patients with edentulous maxillas and mandibles. The trabecular bone pattern of 42 maxillary and 61 mandibular regions of interest (ROIs) was classified into five grades. Grayscale density measurements were made within a marked area of the ROI in the image of the periapical digital radiograph in the same corresponding trabecular region. To obtain conversion values, including the effects of age, gender, and region

an analysis was made to develop regression equations.

*Results* The kappa value for intra- and interobserver differences was 0.71–0.85. The strength of the radiographic conventional value to predict the grayscale density measurement of digital radiography was gained from the regression analysis, with  $R^2 = 0.75$ –0.8. The regression equation for the maxilla and the mandible were separated, and the age, gender, and region of the jaws were included.

*Conclusions* Conventional intraoral radiographic values of the trabecular bone pattern can be converted to values of grayscale density measurements from intraoral digital radiography. The regression equation for the conversion was obtained by including the effects of age, gender, and region of the jaw.

**Keywords** Conversion value - Density measurement - Intraoral conventional radiography - Digital radiography - Regression equation



**Menik Priaminiarti**

**Email:** [menikneidy@yahoo.com](mailto:menikneidy@yahoo.com)

**Email:** [menik-pad@ui.ac.id](mailto:menik-pad@ui.ac.id)

#### References

1. White SC, Pharoah MJ. Oral radiology: principles and interpretation. 5th ed. St. Louis: Mosby; 2004. pp. 71–120, 225–77, 314–29, 677–91.
2. Whaites E. Essentials of dental radiography and radiology. 3rd ed. London: Churchill Livingstone; 2007. p. 289–97.
3. Mupparapu M, Sinner SR. Implant imaging for the dentist. J Can Dent Assoc. 2004;70(1):32.
4. Berkhout WER, Sanderink GCH, van der Stelt PF. Does digital radiography increase the number of intraoral radiographs? A questionnaire study of Dutch dental practices. Dentomaxillofac Radiol. 2003;32:124–7.
5. van der Stelt PF. Filmless imaging: the uses of digital radiography in dental practice. JADA. 2005;136:1379–87.
6. Parks ET, Williamson GF. Digital radiography: an overview. J Contemp Dent Pract. 2002;3(4):23–39.
7. Yoshiura K, Welander U, David WM, Li G, Shi X-Q, Nakayama E, et al. Comparison of the psychophysical properties of various intraoral film and digital systems by means of the perceptibility curve test. Dentomaxillofac Radiol. 2004;33:98–102.
8. Taguchi A, Tanimoto K, Akagawa Y, Wada T, Rohlin M. Trabecular bone pattern of the mandible: comparison of panoramic and CT. Dentomaxillofac Radiol. 1997;26:85–9.
9. Taguchi A, Tanimoto K, Sueti Y, Otani K, Wadamoto M, Akagawa Y, et al. Observer agreement in the assessment of mandibular trabecular bone pattern from panoramic radiographs. Dentomaxillofac Radiol. 1997;26:90–4.
10. Altman DG. Practical statistic for medical research. 1st ed. London: Chapman and Hall; 1991. p. 403–9.

11. Ulm C, Tepper G. Structure of atrophic alveolar bone. In: Georg W, editor. *Implants in qualitatively compromised bone*. London: Quintessence; 2004. p. 29–31.
12. Misch CE. *Contemporary implant dentistry*. 2nd ed. St. Louis: Mosby; 1999. p. 73–118.
13. Nanci A, Whitson SW, Blanco P. Bone. In: Nanci A, editor. *Ten Cate's oral histology, development, structure, and function*. 6th ed. St. Louis: Mosby; 2003. p. 111–44.
14. Neukam FW, Kloss FR. Compromised jawbone quantity and its influence on oral implant placement. In: Zarb G, Lekholm U, Albrektsson T, Tenenbaum H, editors. *Aging, osteoporosis and dental implants*. Chicago: Quintessence; 2002. p. 67–84.
15. Mundy GR. Bone resorbing cells. In: Favus MJ, editor. *Primer on metabolic bone diseases and disorders of mineral metabolism*. 2nd ed. New York: Raven Press; 1993. p. 25–32.
16. Prentice A, Schoenmakers I, Laskey MA, de Bono S, Ginty F, Goldberg GR. Nutrition and bone growth and development. *Proc Nutr Soc*. 2006;65(4):348–60.